

## DIRECTIONS FOR USE

### **Liquid Injection (Furrow Application)**

1. Fill the liquid injection cart with cool, clean, non-chlorinated water.
2. Remove the inoculant pack (calico bag) from the outer plastic bag and suspend the unopened calico bag inside the tank and gently squeeze the bag until the water is discoloured.
3. Agitate or mix the water inside the tank thoroughly for even distribution of the inoculant.
4. Apply at 50-100L/hectare depending on row spacing, as soon as possible.
5. Ensure coarse nozzles (>0.22mm) are used to avoid blockages.

### **Application Using Brushmaster™ Inoculant Applicator**

1. Read the Brushmaster™ applicator operations manual prior to use (available for download from [www.greenmicrobes.com.au](http://www.greenmicrobes.com.au)).
2. Select the applicator nozzle to suit the grain flow rate as per the manual.
3. Ensure that the pump inlet screen is in place.
4. Add cool, clean, non-chlorinated water into the applicator bucket until pump is completely immersed (this will be the baseline for the water).
5. Remove the inoculant pack (calico bag) from the outer plastic bag and place the unopened calico bag into the applicator bucket.
6. Add 2 more litres of water (2 litres for every 1 tonne Inoculeze™ bag).
7. Gently squeeze the bag until the water is discoloured.
8. Switch the disc motor on and ensure that the disc is spinning freely.
9. Switch the pump on and ensure the liquid is flowing freely through the nozzle.
10. Commence grain flow. For best results treated seeds must be sown into a moist soil profile immediately.

**Note:** Thoroughly rinse the applicator and the bucket with clean water after use. Ensure to flush the hose and nozzle with clean water after use.

### **Do's and Don'ts**

Double the application rate of inoculant if the paddock has never grown this pulse previously (i.e. use 2 bags of inoculant without changing the water rate).

Do not mix inoculants with pesticides, fertilisers or other chemicals.

Do not use if sowing into soil with poor moisture (dry sowing) or if the soil temperature is over 30°C.

Do not store opened packs of inoculant.

Do not use out-of-date inoculants.

### **Warnings**

The following conditions may delay or adversely affect nodulation:

- High levels of soil nitrate
- Excessive application of starter nitrogen fertiliser
- Herbicide residues in the soil

This product contains live microorganisms. Avoid contact with eyes, skin rash and abrasions. Avoid inhaling dust. Keep out of reach of children.

Safety Data Sheet available for download from [www.greenmicrobes.com.au](http://www.greenmicrobes.com.au)

### **Warranty**

The product must be used strictly in accordance with the label directions and the instructions. As the manner of the use of this product is beyond the control of the manufacturer, no warranty is given beyond the warranties imposed by the Statutes and in any case, liability is limited to replacement of the product or (at the option of Green Microbes Australia) the refund of the price paid. Buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

\*AIRG approved strain

<sup>†</sup>Viable cell counts at the time of manufacturing.

### **Product of Australia**

Manufactured by:

Green Microbes Australia Pty Ltd.

Unit 5 42 Sam Street

Forbes NSW 2871

Phone: (02) 6851 2788

[contact@greenmicrobes.com.au](mailto:contact@greenmicrobes.com.au)

[www.greenmicrobes.com.au](http://www.greenmicrobes.com.au)